

This Virus Is a Law-Ordered Death Sentence If Your Pet Becomes Infected

It's a fast-moving virus that can start producing signs in only days. The law requires your pet to be quarantined, and if blood tests show he's infected, he must be euthanized. Prevention is the only answer — know what to do to protect your pet to save yourself the worry and potential heartbreak.

Reviewed by [Dr. Becker](#)

STORY AT-A-GLANCE

- The rabies virus is a very serious, usually fatal inflammatory infection affecting the brain and central nervous system
- In the U.S., the rabies virus is most often transmitted to pets through a bite from an infected wild animal
- Once the virus reaches the brain, the infected animal will experience from one to three phases of the infection and associated symptoms
- Diagnosis of rabies in living animals involves a brief quarantine period and fluid samples in symptomatic pets
- Rabies is a fatal disease, but prevention can be achieved with a well-timed initial vaccine against the virus, followed by re-vaccinations as required by law or titer tests for at-risk pets

Editor's Note: This article is a reprint. It was originally published November 12, 2017.

The rabies virus, also called acute viral encephalomyelitis, is an extremely serious, usually fatal inflammatory infection that affects the brain and central nervous system (CNS). Rabies is a single strand RNA virus of the genus *Lyssavirus* in the family *Rhabdoviridae* that's carried in the blood and saliva of infected animals.

How the Rabies Virus Is Transmitted

In dogs and cats in the U.S., the rabies virus is most often transmitted when an infected fox, raccoon, skunk, coyote or bat bites a pet. Infectious virus particles are present in the saliva of these animals to more efficiently transmit the disease.

It's extremely rare, but transmission has also been documented to occur through exposure to the escaping gases of an infected decomposing animal. This transmission usually happens in caves where there are large populations of infected bats. It's something to think about if you have a hunting dog or you like to explore caves with your dog.

Rabies is a zoonotic disease, meaning it can also be transmitted to humans by infected animals. Once the rabies virus enters a pet's body, it multiplies in muscle cells, and then spreads to nearby peripheral sensory and motor nerves, which move it to the brain and central nervous system (CNS).

During the incubation period before the virus enters the CNS, the infected animal doesn't exhibit symptoms and can't transmit the disease. The virus moves relatively slowly, with the average time between exposure to brain involvement being three to eight weeks in dogs, and two to six weeks in cats. However, incubation periods as long as six months have been reported in dogs.

Rabies Phases and Associated Symptoms

Once the rabies virus reaches the brain, the animal will experience one, two or three phases of the infection. Also at this point, the virus moves to the salivary glands where it can be transmitted by a bite.

The first rabies phase is the prodromal phase, which usually lasts between one and two days in cats, and two to three days in dogs. Symptoms during this phase can include a fever, anxiety, nervousness and a desire to be left alone. Normally friendly pets may show shyness, irritability or snappiness.

Normally aggressive animals, interestingly, can turn docile and affectionate. We see that a lot with infected wild animals. Most pets also persistently lick at the site where they were bitten.

The second phase of rabies is either the furious phase or the paralytic phase, which is also called the dumb phase. In the furious phase, which can last from one to seven days, symptoms include restlessness, irritability, hypersensitivity to noise and visual stimuli, roaming and attacking behaviors. Eventually, the animal becomes disoriented, begins having seizures and eventually dies.

The paralytic phase can follow either the prodromal or the furious phase and is usually seen two to four days after the first signs of disease are noticed. Nerves in the head and throat will be affected in this phase, which leads to excessive drooling because the pet can no longer swallow efficiently.

There can also be significant labored breathing, a dropped jaw and choking sounds, as the diaphragm muscles and muscles of the face become paralyzed. The animal will grow progressively weaker and eventually go into respiratory failure and die.

Diagnosing Rabies

Rabies is a fast-moving virus that must be treated as soon as symptoms appear if the pet has any chance for survival. If your dog or cat has been in a fight with another animal, has been bitten or scratched by another animal or if you suspect he may have come into contact with a rabid animal, you should take him to your veterinarian or emergency clinic immediately.

Diagnosis of rabies in a living dog or cat is done through history taking and symptom observation. Pets suspected of having rabies are quarantined in a locked kennel at a veterinary clinic for 10 days and carefully monitored. If your pet is showing progressive symptoms of the disease while quarantined, the veterinarian will take fluid samples for evaluation. If the samples are positive for rabies, sadly, the law requires that your dog or cat be euthanized.

Animals who die of a suspected rabies infection are diagnosed post-mortem (after death) with a direct fluorescence antibody test performed at a state-approved laboratory. All confirmed cases of rabies must be reported to the state health department. Unfortunately, there's no treatment or cure for rabies. Death typically occurs in unprotected animals within seven to 10 days after the onset of symptoms.

Rabies Prevention/Vaccination

Since rabies is a devastating and fatal disease, it's important to protect your dog or cat through vaccination as appropriate and required by law, as well as to avoid situations in which your pet could be exposed to a rabid animal. **Rabies vaccines** are the only vaccines mandated by law in all 50 states.

These vaccines, like all vaccines, have the potential for adverse reactions or reactions that can range from very mild to profound, up to and including anaphylaxis and death. Fortunately, all 50 states now have a three-year revaccination protocol after the second vaccine is given. However, a few county and city laws may differ, so it's important to check with your veterinarian or local animal control agency for more information.

The amount of rabies vaccine given is a hotly debated topic among veterinarians. While the majority of integrative vets believe a "one size fits all" approach to vaccines puts dogs under 40 pounds at risk for substantially more rabies vaccine reactions, the law still mandates all dogs must receive a 1-milliliter dose.

Thankfully, Dr. John Robb has brought this issue to national attention and we are hoping the increased awareness will prompt changes in this requirement on a state-by-state basis.

Some states allow for medical exemptions from rabies vaccines for pets who are ill or have had an adverse reaction in the past.

Additional Recommendations to Improve Vaccination Safety

It's strongly recommended to wait until a pet is as old as allowable by law before giving the first rabies vaccination. In some states, this is six months. In others, it's earlier. A booster shot is usually required within 12 months.

After the second shot, insist on the three-year rabies vaccine. The three-year vaccine is actually identical to the one-year vaccine, so you're providing the protection required by law, but at much less frequent intervals. This means fewer vaccines throughout pets' lifetimes, which means healthier pets.

In addition, rabies vaccinations should never be given in combination with or at the same time as another vaccine. Rabies vaccines should be given alone, separated from all other vaccinations by at least two weeks. Because rabies is the only vaccine required by law, it should be the only repetitive vaccine you give your pet. You can titer for all other diseases.

You can also do a homeopathic detox remedy for the rabies vaccine called Lyssin, especially if your pet has had an adverse reaction. Always closely monitor the injection site after each vaccination. If you notice inflammation or any abnormality or change in the appearance of the skin, such as a lump, irritation or heat, contact your veterinarian immediately.

It's the goal of the integrative veterinary community that the Rabies Challenge Fund, which was initially set up by immunologist Dr. Ron Schultz and veterinarian Dr. Jean Dodds, proves that the rabies vaccine provides duration of immunity for a much longer period than three years.

Ultimately, that would mean pets receive far fewer vaccines. It will allow animals to be immunologically protected without the risk of cumulative vaccine damage over the course of their lifetime.